

Article

Evaluation of a Fixed Spraying System for Phytosanitary Treatments in Heroic Viticulture in North-Eastern Italy

Gaetano Imperatore ¹, Aurora Ghirardelli ^{1,*}, Luca Strinna ¹, Cristiano Baldoin ², Alberto Pozzebon ¹ , Giuseppe Zanin ¹ and Stefan Otto ³

- ¹ Department of Agronomy, Food, Natural Resources, Animals and Environment, DAFNAE, University of Padua, Viale dell'Università 16, 35020 Legnaro, Italy; gaetano.imperatore@unipd.it (G.I.); luca.strinna@unipd.it (L.S.); alberto.pozzebon@unipd.it (A.P.); giuseppe.zanin@unipd.it (G.Z.)
- ² Department of Land, Environment, Agriculture and Forestry, TESAF, University of Padua, Viale dell'Università 16, 35020 Legnaro, Italy; cristiano.baldoin@unipd.it
- ³ Institute for Sustainable Plant Protection (IPSP-CNR), National Research Council of Italy, Viale dell'Università 16, 35020 Legnaro, Italy; stefan.otto@cnr.it
- * Correspondence: aurora.ghirardelli@phd.unipd.it

Abstract: Modern viticulture cannot be practiced without the use of plant protection products to control diseases like downy mildew, powdery mildew, and pests. In severely sloping vineyards, where mechanization is not possible, pesticide application is realized using spray guns, which is a laborious, expensive, and dangerous application technique. In these vineyards, where viticulture is defined as “heroic viticulture,” vine-growers could seriously take advantage of innovation in spray-technique applications. For this reason, several prototypes of a fixed spraying system (FSS) were realized in recent years. Two prototypes of a fixed spraying system were built in 2019 in two different vineyards in the Veneto region (north-eastern Italy). In both vineyards, the fixed spraying systems were used to perform pesticide application during the 2020 season to control downy mildew, powdery mildew, and pests. With this solution, both vineyards were successfully protected, resulting in comparable infection degrees and yields as the ones protected with airblast sprayers and spray guns. This study contributes to assert fixed spraying systems as an innovation that could improve working conditions, safety, timing, and performances of plant protection products’ application in heroic viticulture areas.

Keywords: fixed spraying system; steep-slope viticulture; pesticide application; downy mildew; organic viticulture; solid set canopy distribution system



Citation: Imperatore, G.; Ghirardelli, A.; Strinna, L.; Baldoin, C.; Pozzebon, A.; Zanin, G.; Otto, S. Evaluation of a Fixed Spraying System for Phytosanitary Treatments in Heroic Viticulture in North-Eastern Italy. *Agriculture* **2021**, *11*, 833. <https://doi.org/10.3390/agriculture11090833>

Academic Editor: Jin He

Received: 27 July 2021

Accepted: 28 August 2021

Published: 31 August 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Hilly and mountain viticulture present greater difficulties and costs of cultivation than plain viticulture. In Italy, according to the last census of agriculture, the area planted with vineyards in hilly and mountainous areas amounts to 357,000 and 35,800 ha, respectively, representing about 64% of the national area. In most of these vineyards, the cultivation is mechanized, and the production costs are just slightly higher than plain viticulture ones. The situation is different when it comes to viticulture practiced on severe sloping hills or in mountainous areas. A specific name has been given for this type of viticulture by the Centre for Research, Environmental Sustainability, and Advancement of Mountain Viticulture [1]: heroic viticulture. This term applies to vine growing occurring on slopes greater than 30%, in vineyards above 500 m a.s.l. (excluding highlands), or in vineyards on terraces. In such contexts, the cultivation is hardly mechanizable, and consequently, the production cost of the grape is higher, as the majority of growing activities are carried out by hand following traditional agricultural practices.

Commercial vineyards often require intensive crop protection techniques with several pesticide applications. Fungicide treatments against downy mildew, powdery mildew, and

botrytis, and insecticide applications against the leafhopper *Scaphoideus titanus* Ball, vector of flavescence dorée, and the grape berry moth *Lobesia botrana* (Denis & Schiffermüller), are the most common treatments.

In heroic vineyards treatments are carried out entirely by hand, using a knapsack mistblower or high-volume sprayer equipped with a spray gun [2] (hereafter: SG). The use of SG in terraced vineyards is laborious and time-consuming. Moreover, since inter-rows are very narrow (no more than 1.5 m) and the operating pressure often exceeds 20 bars [3], this application technique puts the operator into direct contact with the applied solution, resulting in a high risk of pesticide exposure for operators.

Despite the high volumes and high pressures involved, the biological efficiency of the treatments show remarkable variability related to operator precision: some parts of the canopy could be improperly sprayed, while in others, there may be an over-dosage and dripping to the soil of the excess solution [4].

Nevertheless, a fixed spray system (hereafter: FSS) can be suggested also in vineyards where air-blast sprayers (hereafter: ABS) are used, for the protection of difficult rows, like those at the hedge of fields where ABS can hardly operate.

In heroic viticulture, crop protection is one of the most time-consuming activities. According to the estimates of the Prosecco DOCG (*Denominazione di Origine Controllata e Garantita*) Consortium of Conegliano and Valdobbiadene [5], a single pesticide application in severely sloping vineyards requires 4–6 h/ha of work, whereas the same task in plain and highly mechanized plain viticulture requires 0.75–1 h/ha. Improving time-effectiveness could help increase the income of farmers, resulting in better preservation of the viticultural landscape. In severely sloping vineyards, crop protection is also the most cost-intensive operation. In steep slope viticulture, most of the field operations can be done exclusively manually, resulting in a cost difference of +161% compared to plain viticulture [6].

FSSs have already been studied and applied in fruit cultivation, in particular for the control of fungal pathogens in apple [7–9] and cherry orchards [10] as well as for controlling pests [11]. However, most of the previous prototypes used in orchards are not suitable for vineyards, because they are designed to use the existent anti-frost irrigation system, for the distribution of calcium polysulfide solutions [12]. This technique works against the apple scab, but it is not effective against the typical pathogens of the vine, such as downy mildew and powdery mildew, which require the covering of the bunches. For this reason, other types of FSSs have been designed specifically for viticulture. Some prototypes of FSS were built and tested recently in Italy [13–15] and in the USA [16,17], which also have off-field drift potential [13].

Therefore, the FSS for heroic viticulture treatments, rather than being an adaptation of the irrigation system, must be specifically designed to apply the plant protection product as a sprayer and cover the crop canopy with droplets of appropriate number and size, with acceptable off-field drift.

The aim of the study was to compare the biological performance of FSS prototypes with the farmer standard spray-application techniques in two peculiar areas of the Veneto region: the Prosecco DOCG Consortium area of Conegliano and Valdobbiadene, a United Nations Educational, Scientific, and Cultural Organization World Heritage Site [18], and the Soave Consortium area, recognized by the Food and Agriculture Organization as a Globally Important Agricultural Heritage Site [19]. The two vineyards selected for the study present different training systems, vine varieties, crop protection methods (conventional and organic agriculture), and geological and meteorological conditions. Therefore, the FSS was designed and tested to meet the specific needs of two different agronomical contexts.

2. Materials and Methods

2.1. Experimental Areas

Experimental trials were performed in two different vineyards. The first vineyard was in Vidor (Treviso, Italy 45°53'9.29" N, 12°3'18.59" E, 147 m a.s.l., Scandolera Wineries), within the Prosecco DOCG Consortium area. Vines were in linear trellis, and crop protection

was conventional. The FSS was mounted in a steep vineyard (slope 35%) of the Glera vine variety, the principal vine of Prosecco DOCG wine. The area chosen for the trial was a 750 m² vineyard where vines are trained to the local traditional “doppio capovolto” (Figure 1a) linear training system. The spacing of the vines was not regular, ranging from 2.50 × 1.00 m to 3.20 × 1.40 m, due to irregular terracing soil conformation. The age of the plants varied from 2 to 40+ years old, but most of them were 15/20 years old. The area chosen for the trial had six rows, each one 45-m long, plus a 50-m long row for the untreated control plot (hereafter: UTC).



(a)

(b)

Figure 1. Training systems in experimental vineyards: (a) Vidor “Doppio capovolto” training systems, (b) Soave “Pergola veronese” training system.

The second vineyard was in Soave (Verona, Italy 45°27'45.39" N, 11°13'59.40" E, 347 m a.s.l. Coffele vineries), within the Soave DOC Consortium area. Vines were in Pergola trellis, and crop protection was according to organic farming protocol. The area chosen for the trial was a plain 2000-m² wide, organic vineyard. The FSS was mounted on a vineyard of the Garganega vine variety trained to the local traditional “Pergola Veronese” (Figure 1b) trellis training system with vines spaced 3.50 × 1.00 m. The area chosen for the trial had six rows, each one 85-m long, plus a 60-m long row for UTC.

2.2. Experimental Design

In Vidor, the FSS was compared with SG. The experimental area was split into two plots: each one of the six rows had half of its length treated with the FSS, and the other half treated with a SG (Figure 2a). This splitting without plot randomization was mandatory due to technical-constructive reasons related to the FSS structure. A UTC was included.

In Soave, the FSS was compared with ABS. For the same reasons as above, also in Soave, the experimental area was split into two non-randomized plots (Figure 2b). A UTC was included.




Figure 2. Experimental design in the Vidor vineyard (a) and in the Soave vineyard (b).

2.3. Fixed Spraying Systems

The FSS was designed taking into account the results of previous experiments carried out on vineyards [14] and orchards [7,20,21]. According to these findings, the general layout of the plant included a lower line and an upper line arranged to spray separately the fruit zone and the upper canopy, to ensure good coverage of the leaves since the most relevant pathogens were located on the downsides (e.g., downy mildew). For this purpose, the general layout of the FSS was designed as follows (Figure 3).




Figure 3. Scheme of the fixed spray system.

A polyethylene tank, with a 1-m³ capacity, was placed at the head of the vineyard and equipped with an electric-driven three-membrane pump, with a 40-L·min⁻¹ capacity; a general supply tubing Ø 25 mm connected the tank with a manifold from which the pipes direct to the individual rows branch off. The piping that feeds each row was in turn divided into two separate lines, one for the spraying of the foliage from above, the other for the clusters, and the lower part of the vegetation from below. Due to the different locations of the two lines, they were fitted with different types of nozzles, each one provided with an anti-drop valve calibrated to open when the pressure exceeded 2 bar: the lower line was equipped with TeeJet TXR800053VK hollow cone nozzles, with a flow rate of 0.21 L/min @ 3 bar, oriented 45° upwards to target the fruit zone and the lower part of the canopy; the upper line, installed about 50 cm over the top of the canopy, holds micro-sprinklers spraying downwards. The specific components differ between the two versions of the plant installed in the experimental vineyards and will be described in detail at the end of this section. At the end of the row, the tubes converge into a transverse pipe, which returns

to the tank, thus reproducing the functionality of a normal hydraulic closed circuit, like the ones that are normally used on the tractor-driven sprayers.

The whole tubing was also fed with compressed air with the function of emptying the lines and the nozzles after spraying. All pipes were supplied individually and isolated from the others through solenoid valves placed at both the ends of the individual row pipes; the operation sequence of the valves was controlled by a programmable logic controller (hereafter: PLC).

The operational steps of the spraying process are:

1. the agrochemical mixture is circulated at low pressure in the system, with the solenoid valves of the row to be treated open, until the liquid completely fills the line; this step takes about 100 s;
2. once the pipeline is fully replenished, the PLC closes the solenoid valve located at the end of the row; the pressure rises suddenly at 6 bar, the anti-drop valves open, and the nozzles start spraying, with virtually no delay between the opening of the first and last nozzle; after 15 s, the PLC opens the terminal row valve: the pressure drops immediately below 2 bar, anti-drop valves close, and the mixture flows through the collecting tube back to the tank;
3. at the end of spraying, the pump is switched off and a solenoid valve lets compressed airflow at low pressure (<0.2 bar) into the supply line, thus emptying the mixture into the tank; after 70–80 s, the air pressure in the nozzle line increases due to the closing of the end-of-row valve, and the residual mixture in the nozzles is sprayed on the foliage;
4. steps 2 and 3 are repeated for each row.

2.4. Application of Pesticides

The application of pesticides in Vidor was carried out using the FSS or SG. The use of the SG is the traditional application technique of pesticide in the Prosecco DOCG area because the majority of vineyards there are planted on terraces in sloping hills. The application of pesticide carried out with the FSS and the one performed with the hand SG were both set to deliver a 1000 L/ha volume of phytosanitary solution, in line with the mean volumes used in the area for pesticide application on vineyards. The SG method was set to spray at 20 atm of pressure, while the FSS was set to spray at 6 atm of pressure in the lower line and 3.5 atm in the upper line. The details of pesticide applications are listed in Tables 1 and 2.

Table 1. Dates, product names, rates, and active ingredients, and target of plant health treatments realized in the Vidor vineyard during 2020.

Date	Product Name	Rate/ha	Active Ingredient	Target
2 May	Polyram DF	1500 g	Metiram	Downy and powdery mildew
	Dimethomorph	400 mL	Dimethomorph	Downy and powdery mildew
	Sulfur	3000 mL	Sulfur	Powdery mildew
14 May	Polyram DF	1500 g	Metiram	Downy and powdery mildew
	Dimethomorph	400 mL	Dimethomorph	Downy and powdery mildew
	Sulfur	3000 mL	Sulfur	Powdery mildew
21 May	Sercadis	150 mL	Fluxapyroxad	Powdery mildew
	Enervin TOP	2500 g	Ametoctradin + metiram	Downy mildew
1 June	Sercadis	150 mL	Fluxapyroxad	Powdery mildew
	Zorvec Zelavin	1500 mL	Oxathiapiprolin	Downy mildew
12 June	Lieto SC	450 g	Cymoxanil + zoxamide	Downy mildew
	Quantum	500 mL	Dimethomorph	Downy and powdery mildew
	Score 25 EC	200 mL	Difenoconazole	Powdery mildew and black rot

Table 1. Cont.

Date	Product Name	Rate/ha	Active Ingredient	Target
23 June	Enervin Duo	1000 mL	Ametoctradin + dimethomorph	Downy mildew
	Vivando	200 mL	Metrafenone	Powdery mildew
	Kestrel	400 mL	Acetamiprid	Leafhoppers
6 July	Lieto SC	450 g	Cymoxanil + zoxamide	Downy mildew
	Vivando	200 mL	Metrafenone	Powdery mildew
14 July	Ridomil GOLD R WG	5000 g	Metalaxyl + copper oxychloride	Powdery mildew
	Zorvec Zelavin	300 mL	Oxathiapiprolin	Downy mildew
	Brionflo 100 SC	1000 mL	Cyazofamid	Downy mildew
	Kestrel	400 mL	Acetamiprid	Leafhoppers
21 July	Ridomil GOLD R WG	5000g	Metalaxyl + copper oxychloride	Powdery mildew
	Champ 20	1000 g	Copper hydroxide	Downy mildew
	Sulfur	3000 mL	Sulfur	Powdery mildew
30 July	Curame bordeaux	5000 g	Copper sulfate + cymoxanil	Downy mildew
	Mavrik smart	300 mL	Tau-fluvalinate	Leafhoppers
	Zolvis	4000 g	Sulfur	Powdery mildew

Table 2. Operating parameters used in every pesticide application in the Vidor vineyard during 2020.

Thesis	Nozzle Type	Number of Nozzles	Operating Pressure (bar)	Air Pressure (bar)	Flow Rate @ 3.0 bar (L min ⁻¹)	Forward Speed (km h ⁻¹)	Volume Rate (L ha ⁻¹)
FSS	Upper line	Netafim Coolnet Pro mod. 075	1 nozzle every 0.8 m	3.5	8	0.43	-
	Lower line	TeeJet TXR TXR800053VK (hollow cone)	A couple every 1.2 m	5–5.5	8	0.21	1000
SG	-	TeeJet TXA8004VK (hollow cone)	1	20	-	1.58	2.5

The application of pesticides in Soave was carried out using the FSS or ABS (Florida NAZA 600 L at 25 bar), both set to deliver a 750 L/ha volume of phytosanitary solution, in line with the mean volumes used in the area for pesticide application on vineyards. The details of pesticide applications are listed in Tables 3 and 4. Since in organic farming most of the active substances are forbidden, only copper- and sulfur-based fungicides, and only authorized insecticides, were used during the field trial, according to Commission Regulation (EC) No 889/2008 [22].

Table 3. Dates, product names, rates, active ingredients, and the target of plant health treatments realized in the Soave vineyard during 2020.

Date	Product Name	Active Ingredient	Rate/ha	Target
2 May	Mexiram Hi Bio	Copper hydroxide	1.5 Kg/ha	Downy mildew
	Tiovit Jet	Sulfur	3 kg/ha	Powdery mildew
9 May	Mexiram Hi Bio	Copper hydroxide	1.5 Kg/ha	Downy mildew
	Tiovit Jet	Sulfur	3 kg/ha	Powdery mildew
14 May	Mexiram Hi Bio	Copper hydroxide	1.5 Kg/ha	Downy mildew
	Tiovit Jet	Sulfur	3 kg/ha	Powdery mildew
21 May	Mexiram Hi Bio	Copper hydroxide	1 Kg/ha	Downy mildew
29 May	Tiovit Jet	Sulfur	3 kg/ha	Powdery mildew
	Mexiram Hi Bio	Copper hydroxide	1.5 Kg/ha	Downy mildew

Table 3. Cont.

Date	Product Name	Active Ingredient	Rate/ha	Target
3 June	Tiovit Jet	Sulfur	3 kg/ha	Powdery mildew
	Mexiram Hi Bio	Copper hydroxide	1.5 Kg/ha	Downy mildew
5 June	Tiovit Jet	Sulfur	3 kg/ha	Powdery mildew
	Mexiram Hi Bio	Copper hydroxide	1.5 Kg/ha	Downy mildew
8 June	Tiovit Jet	Sulfur	3 kg/ha	Powdery mildew
	Mexiram Hi Bio	Copper hydroxide	1.5 Kg/ha	Downy mildew
15 June	Tiovit Jet	Sulfur	3 kg/ha	Powdery mildew
	Mexiram Hi Bio	Copper hydroxide	1.5 Kg/ha	Downy mildew
25 June	Microthiol Disperss	Sulfur	4 kg/ha	Powdery mildew
	Mexiram Hi Bio	Copper hydroxide	0.5 kg/ha	Downy mildew
7 July	Microthiol Disperss	Sulfur	6 kg/ha	Powdery mildew
	Neemik Ten	Azadirachtin	2.6 L/ha	Leafhoppers
15 July	Microthiol Disperss	Sulfur	8 kg/ha	Powdery mildew
28 July	Mexiram Hi Bio	Copper hydroxide	400 g/hl	Downy mildew
	Microthiol Disperss	Sulfur	800 g/hl	Powdery mildew
18 August	Poltiglia Disperss	Bordeaux mixture	2 kg/ha	Downy mildew
	Microthiol Disperss	Sulfur	4 kg/ha	Powdery mildew
	Surround Wp	Kaolin	15 kg/ha	

Table 4. Operating parameters used in every pesticide application in the Soave vineyard during 2020.

Thesis	Nozzle Type	Number of Nozzles	Operating Pressure (bar)	Air Pressure (bar)	Flow Rate @ 3.0 bar (L min ⁻¹)	Forward Speed (km h ⁻¹)	Volume Rate (L ha ⁻¹)
FSS	Upper line	NETAFIM Gyronet 58 SR	1 nozzle every 2.5 m	6	10	0.83	-
	Lower line	TeeJet TXR TXR800053VK (hollow cone)	A couple every 1 m	6	10	0.21	750
ABS	-	TeeJet TXB8003VK (hollow cone)	6	20	-	1.18	5.3

2.5. Meteorological Data

Figure 4 shows the mean daily temperature and precipitation in the two vineyards in the 2020 growing season recorded at meteorological stations located inside or very near the vineyards. In Vidor, the total amount of precipitation between 1 April 2020 and 30 September 2020 was 916.6 mm, and the number of rainy days was 59 (a rainy day was considered a day when the amount of rainfall was over 1 mm). In Soave, the total amount of precipitation in the same period was 546.2 mm, and the number of rainy days was 45.

2.6. Evaluation of Disease Symptoms and Arthropods Abundance

In 2020, the phytosanitary status of leaves and bunches was monitored in both sites, and disease symptoms were evaluated for the spray-application techniques under comparison (FSS, SG, and ABS) and for UTC.

In Vidor, the damage from downy mildew was assessed first on leaves at the time of veraison (BBCH 80; 30 July 2020) and then on bunches at the time of harvest (11 September 2020). In Soave, the evaluation of the leaf damage was assessed at the time of veraison, and the damage on bunches was assessed at the time of harvest (26 September 2020).




Figure 4. Mean daily temperature and precipitation in the Vidor (a) and Soave (b) vineyards in 2020.

To evaluate the performance of the pesticides, the response of the plants to downy mildew disease was analyzed regularly at each growth stage (BBCH-identification keys of grapevine), following the developmental scale described by Lorenz et al. (1994). We scored 100 leaves and 100 bunches picked randomly from 10 central plants of each plot and estimated the percentage of affected organs (disease incidence) and the percentage areas of leaves and bunches showing disease symptoms (disease severity). For the evaluation of the damage in each site, 50 leaves/bunches were sampled randomly from central plants of each plot. The first and the last rows of the vineyards were excluded from sampling to create a buffer zone between the experimental areas and the surrounding vineyards. In Vidor, four plots for FSS and SG, and the UTC, were sampled. In Soave, due to the different arrangement of the FSS, three plots for FSS and ABS, and the UTC, were sampled. Disease severity (infection degree (ID)) was calculated using a scale of eight classes (0–7) with the Townsend–Heuberger formula [23]:

$$ID\% = \frac{\sum n \times v}{N \times V} \times 100, \quad (1)$$

with n = number of leaves/bunches per each class, v = value of the class, N = number of leaves/bunches sampled, and V = value of the higher class of the scale.

The percentage incidence of disease (I_c) was calculated as the percentage of total organs sampled showing symptoms of disease.

In addition to fungal pathogens, mites and leafhoppers occurring in vineyards were also monitored. The abundance of the predatory mite *Kampimodromus aberrans* (Oudemans) and its prey the mite pest *Eotetranychus carpini* (Oudemans) were quantified on 100 leaves per the spray-application techniques and collected in every plot in both vineyards. The leaves were then observed under the stereoscope, and the mite numbers were quantified. Samples were taken at 10-day intervals between 11 June and 28 August in Vidor, while, in Soave, the samples were taken from 19 June to 24 September.

To quantify the abundance of the leafhopper pests *Scaphoideus titanus*, *Zygina rhamni* (Ferrari), *Erasmuneora vulnerata* (Fitch), and *Empoasca vitis* (Goethe), a yellow sticky trap (12 × 20 cm) was placed in every plot in both vineyards. The traps were replaced regularly, and the leafhoppers captured were quantified. In Vidor, the collection of traps was accomplished every ten days from 2 July to 28 August, while, in Soave, the collection was done every 30 days from 10 July to 12 September.

The quantification of the yields was performed in a similar way in the two sites. In Vidor, 10 plants per plot were chosen randomly, and the bunches were divided into two classes: first quality bunches (damage classes 1–2) and second quality bunches. In Soave, since the vineyards are fairly bigger than in Vidor, 20 plants per plot were harvested. Given the heterogeneous nature of the plants and the presence of failed plants, we decided to determine the yield as “kilograms of grapes per vine.”

2.7. Data Analysis

The non-parametric one-factor analysis (Kruskal–Wallis test; post hoc: Conover test; $\alpha = 0.05$) was used to evaluate significant differences in disease severity (Ic) between the spray-application techniques and the UTC in both vineyards.

Data on the abundance of leafhoppers in Vidor collected before the first insecticide application (July 12) was analyzed using the non-parametric one-factor analysis (Kruskal–Wallis test; $\alpha = 0.05$) to evaluate the abundance of leafhopper populations before the insecticide application. Data collected after the first insecticide application were analyzed with the Friedman two-way non-parametric repeated measure analysis of variance ($\alpha = 0.05$) to test the effect of insecticide application using different spray-application techniques on leafhopper populations. Data on the abundance of leafhoppers observed in Soave and collected after insecticide application were analyzed using a Friedman two-way non-parametric repeated measure analysis of variance ($\alpha = 0.05$). The populations of leafhoppers in UTC in both vineyards were not assessed, because insecticide treatments in the Veneto region are mandatory to control grapevine flavescente dorée and its vector *Scaphoideus titanus* [24]. Different analyses were performed on different leafhopper species. In all analyses, the numbers of leafhoppers belonging to the different species observed on each sticky trap were considered as dependent variables (with repeated measures made at different sampling dates, in case of post-treatment sampling), while the spray-application technique was considered as an independent variable.

The effect of pesticide application using different spray-application techniques on mite populations was evaluated using a Friedman two-way non-parametric repeated measure analysis of variance ($\alpha = 0.05$) on the number of predatory and pest mites observed in the two vineyards. Different analyses were performed on different mite species. In all analyses, the numbers of mites per leaf belonging to the different species were considered as dependent variables with repeated measures made at different sampling dates, while the spray application technique was considered an independent one. The non-parametric one-factor analysis (Kruskal–Wallis test; post hoc: Conover test; $\alpha = 0.05$) was used to evaluate significant differences in yield between the spray-application techniques in both vineyards, analyzing separately 1st quality bunches from 2nd quality bunches.

3. Results

3.1. Vidor

The incidence and severity obtained by sampling the four plots for each spray application technique were analyzed with the Kruskal–Wallis test. The comparison was made between FSS, SG, and UTC. The results in Table 5 show a significantly different incidence (Ic) on SG, compared to FSS and UTC. The analysis of severity (ID) also showed significant differences between the three application techniques.

Table 5. Infection of grapevines with downy mildew in the Vidor vineyard: comparison between the two spray-application techniques and the untreated control. Damage index on 23 July 2020 was applied to leaves and, on 11 September 2020, was applied to bunches. Different letters indicate significant statistical differences. * = p -value < 0.05; ** = p -value < 0.01.

Pathogen	Evaluation Day	Application Method	Incidence (ID) (%)	Severity (Ic) (%)
Downy mildew	23 July 2020	Fixed spray system	25.5 a	9.07 b
		Spray gun	13.00 b	3.07 a
		Untreated control	42.47 a	29.80 c
		p -value	0.0181 *	0.0072 **
Downy mildew	11 September 2020	Fixed spray system	4.00 a	1.35 b
		Spray gun	4.00 a	0.85 a
		Untreated control	44.00 b	21.00 c
		p -value	0.0053 **	0.0089 **

The results of the incidence and severity on bunches resembled those on leaves. The incidence and severity were significantly different according to the Kruskal–Wallis test. Incidences on FSS and SG were not significantly different, while UTC had a significantly higher percentage of affected bunches. The severity results replicated the situation on the leaves, distinguishing significant differences between all three application techniques (Table 5).

The yield of 1st quality bunches (CI = 1–2) showed no significant differences between FSS and SG, settling around 9 kg/plant. The yield of 2nd quality bunches (CI >= 3) showed significant differences between the two spray methods, with a lower quantity of 2nd quality bunches in SG than in FSS (Figure 5a).




Figure 5. Grape yield in Vidor (a) and Soave (b) vineyards calculated as kg/plant of vines harvested. Class 1–2 bunches were 1st quality bunches (portion of bunch damaged < 5%); class ≥ 3 bunches were 2nd quality bunches (portion of bunch damaged ≥ 5%). Different letters within each class indicate significant statistical differences.

In this vineyard, the presence of the leafhoppers *S. titanus*, *Z. rhamni*, and *E. vitis* and the mites *K. aberrans* and *E. carpini* were observed (data not shown). The Kruskal–Wallis test on the population of *Z. rhamni* (p -value = 0.546) and on the population of *E. vitis* (p -value = 0.090) on June 22 showed no significant differences between FSS and SG. Just one *S. titanus* was captured during 2020 (on 2 August), so the analysis was not performed for this species. The post-treatment Friedman test on the population of *Z. rhamni* (p -value = 0.179) and on the population of *E. vitis* (p -value = 0.654) showed no significant differences between FSS and SG. The presence of the mite pest *E. carpini* was observed at a low level. However, the Friedman test revealed that the *E. carpini* density was slightly higher in SG than FSS (p -value = 0.014). No differences between spray systems emerged on *K. aberrans* populations (p -value = 0.414).

3.2. Soave

Nearly no leaf damage was detected for most of the season on every thesis. On the other hand, some damage was detected on bunches due to powdery mildew and grey mold after veraison. As for Vidor’s vineyard, the analysis of the incidence data (Ic) of powdery mildew on bunches showed significant differences between theses: FSS was less affected than ABS and UTC. The Kruskal–Wallis test performed on the ID showed no differences between the spray methods. Table 6 shows that the average level of infection at harvest (26 September 2020) was fairly low.

Table 6. Infection of grapevines with powdery mildew and grey mildew in Soave vineyard: comparison between the two spray-application techniques and the untreated control. Different letters indicate significant statistical differences. * = p -value < 0.05.

Pathogen	Evaluation Day	Application Method	Incidence (ID) (%)	Severity (Ic) (%)
Powdery mildew	26 September 2020	Fixed spray system	4.00 a	2.85 a
		Airblast sprayer	6.66 b	3.04 a
		Untreated control	6.67 b	3.61 a
		p -value	0.0456 *	0.8068
Grey mildew	26 September 2020	Fixed spray system	27.33 a	10.38 a
		Airblast sprayer	34.04 a	13.06 a
		Untreated control	32.00 a	16.28 a
		p -value	0.3932	0.2019

The Kruskal–Wallis test on the incidence (Ic) and severity (ID) of botrytis symptoms on bunches showed no differences between FSS, ABS, and UTC at the time of harvest (Table 6).

The yield of 1st quality bunches (CI = 1–2) showed no significant differences between ABS and FSS, with a yield of 8.05 kg/plant for ABS and a yield of 7.21 kg/plant for FSS. The yield of 2nd quality bunches (CI >= 3) showed no significant differences between the two spray methods (Figure 5b).

Leafhoppers *E. vulnerata*, *S. titanus*, *Z. rhamni*, and *E. vitis*, and the predatory mite *K. aberrans*, were observed in this vineyard (data not shown). The Friedman test found no significant differences in leafhopper populations between FSS and ABS (p -value = 0.533). No differences between spray systems emerged on *K. aberrans* populations (p -value = 0.654).

4. Discussion

During 2020, different pest management strategies showed similar results in both vineyards equipped with the FSS.

In Vidor, the major disease was downy mildew. On leaves, the incidence of downy mildew was higher in FSS and UTC than in SG. However, considering the severity of the damage (Ic), it can be seen that the disease was present in several plants of the FSS, but the damage was not as severe as in the UTC (Table 5). This is the reason why the damage on

leaves appears to be less critical in FSS than in UTC. Focusing on bunches' damage, the FSS provided efficient protection against diseases, despite the fact that the SG had a higher canopy penetration potential thanks to a higher operational pressure (6 bar and 20 bar, respectively). Several previous studies on FSSs and solid set canopy delivery systems (SSCDS) determined a lower canopy penetration capability of FSSs and SSCDS than ABS, in particular regarding the lower surface of leaves [25,26]. We empirically observed that the FSS had a lower capability of reaching the lower surface of the leaves, and this can explain the higher incidence of downy mildew on leaves (Table 5). However, the use of systemic active ingredients, and the position of bunches, which are easier to be reached from the lower line of nozzles, allowed achievement of the same ID (%) and yield as in SG.

In Soave, the damage on leaves and bunches due to fungal infections were quite low, even if only active ingredients allowed in organic agriculture were used. The bunches damage index (ID%) showed that the FSS was able to manage the infection of powdery mildew better than the ABS (Table 6). However, the damage caused to bunches by two severe hailstorms, which both occurred after the end of the treatments (on 15 August and 2 September), led to a severe late grey mildew infection. This infection caused so much damage to the UTC that no bunches were spared at the moment of harvesting. It is likely that this high level of damage was related to the plot position at the edge of the vineyard, which was much more exposed to the hailstorm than other plots.

No differences among spray systems were found on arthropods occurring in vineyards highlighting that FSSs are efficient as standard methods for insecticide application against leafhoppers, which are major grapevine pests [27–29], and no differences on side-effects on beneficial predatory mites emerged [30].

5. Conclusions

After a season-long study, FSS biological performances were comparable to those of commercial spraying application techniques in both sites. However, to achieve those results, it is necessary to perform accurate pruning in the early stages of growing, possibly twice in sturdy vines. The financial feasibility is, instead, doubtful, mainly because expensive nozzles for air-blast sprayers were used. At the moment, this makes the cost of the prototype too high for standard commercial vine-growers, even if it can be markedly reduced by involving industrial stakeholders, mainly for the production of nozzles specific for FSSs.

Further research is needed to confirm the biological and hydraulic efficiency of a FSS over years. This will also help to evaluate the economic feasibility of the FSS and the chance of this new technology gaining importance in the market of agricultural sprayer systems.

Author Contributions: Conceptualization, S.O. and G.I.; methodology, G.Z. and A.P.; validation, S.O., G.Z. and A.P.; formal analysis, L.S.; investigation, G.I. and C.B.; writing—original draft preparation, G.I.; writing—review and editing, A.G., S.O. and C.B.; visualization, L.S.; supervision, G.Z.; project administration, G.I.; funding acquisition, G.I. and S.O. All authors have read and agreed to the published version of the manuscript.

Funding: This work was supported by the Rural Development Plan for the Region of Veneto for 2014–20 (Veneto RDP), Measure 16.1.1 and 16.1.2 (grant number: 3685655).

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: The data presented in this study are available on request from the corresponding author.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. CERVIM 2021. Available online: <http://www.cervim.org/en/mountain-and-steep-slope-viticulture.aspx> (accessed on 28 August 2021).
2. Michael, C.; Gil, E.; Gallart, M.; Stavrinides, M.C. Evaluation of the effects of spray technology and volume rate on the control of grape berry moth in mountain viticulture. *Agriculture* **2021**, *11*, 178. [[CrossRef](#)]
3. Dubuis, P.-H.; Monod, V.; Gindro, K. Pulvérisateurs de type gun et canon: Étude de littérature sur leur utilisation et les risques spécifiques. *Rev. Suisse Vitic. Arboric. Hortic.* **2019**, *51*, 300–305.
4. Michael, C.; Gil, E.; Gallart, M.; Stavrinides, M.C. Influence of spray technology and application rate on leaf deposit and ground losses in mountain viticulture. *Agriculture* **2020**, *10*, 615. [[CrossRef](#)]
5. Merlo, R. Consorzio di Tutela del Conegliano Valdobbiadene Prosecco Superiore DOCG. Personal communication, 2021.
6. Strub, L.; Loose, S.M. The cost disadvantage of steep slope viticulture and strategies for its preservation. *Oeno One* **2021**, *55*, 49–68. [[CrossRef](#)]
7. Landers, A.; Agnello, A.; Shayya, W. Engineering challenges in the development of a fixed spraying system for high-density apple trees. In Proceedings of the III workshop on Spray Application Techniques—Fruit Growing, Barcelona, Spain, 9–12 July 2005.
8. Verpont, F.; Favareille, J.; Zavagli, F. Fixed spraying system: A future potential way to apply pesticides in an apple orchard? In Proceedings of the Workshop on Spray Application in Fruit Growing, Lake Constance, Germany, 15–18 July 2015; pp. 53–54.
9. Sinha, R.; Ranjan, R.; Bahlol, H.Y.; Khot, L.R.; Hoheisel, G.A.; Grieshop, M.J. Development and performance evaluation of a pneumatic solid set canopy delivery system for high-density apple orchards. *Trans. ASABE* **2020**, *63*, 37–48. [[CrossRef](#)]
10. Niemann, S.M.; Whiting, M.D. Spray coverage in apple and cherry orchards using a solid set canopy delivery system. *Acta Hort.* **2016**, *1130*, 647–653. [[CrossRef](#)]
11. Owen-Smith, P.; Wise, J.; Grieshop, M.J. Season long pest management efficacy and spray characteristics of a solid set canopy delivery system in high density apples. *Insects* **2019**, *10*, 193. [[CrossRef](#)] [[PubMed](#)]
12. Bondesan, D.; Rizzi, C.; Dorigoni, A.; Pasqualini, J.; McHeli, F. First assessments of fixed spray application systems in narrow-wall apple orchards. *Asp. Appl. Biol.* **2016**, *132*, 411–414.
13. Otto, S.; Loddo, D.; Schmid, A.; Roschatt, C.; Venturelli, M.; Innerebner, G. Droplets deposition pattern from a prototype of a fixed spraying system in a sloping vineyard. *Sci. Total Environ.* **2018**, *639*, 92–99. [[CrossRef](#)]
14. Baldoïn, C.; Causin, R.; Loddo, D.; Otto, S.; Duso, C.; Zanin, G. Design and first testing of a prototype of a fixed spraying system for sloping vineyards. *Asp. Appl. Biol.* **2016**, *132*, 397–404.
15. Innerebner, G.; Roschatt, C.; Schmid, A. Efficacy of fungicide treatments on grapevines using a fixed spraying system. *Crop Prot.* **2020**, *138*, 105324. [[CrossRef](#)]
16. Sinha, R.; Ranjan, R.; Khot, L.R.; Hoheisel, G.A.; Grieshop, M.J. Comparison of within canopy deposition for a solid set canopy delivery system (SSCDS) and an axial-fan airblast sprayer in a vineyard. *Crop Prot.* **2020**, *132*, 105124. [[CrossRef](#)]
17. Sinha, R.; Khot, L.R.; Hoheisel, G.A.; Grieshop, M.J.; Bahlol, H. Feasibility of a Solid set canopy delivery system for efficient agrochemical delivery in vertical shoot position trained vineyards. *Biosyst. Eng.* **2019**, *179*, 59–70. [[CrossRef](#)]
18. UNESCO Le Colline del Prosecco di Conegliano e Valdobbiadene. Available online: <https://whc.unesco.org/en/list/1571/> (accessed on 28 August 2021).
19. FAO Soave Traditional Vineyards, Italy. Available online: <http://www.fao.org/giahs/giahsaroundtheworld/designated-sites/europe-and-central-asia/soave-traditional-vineyards/en/> (accessed on 28 August 2021).
20. Agnello, A.; Landers, A.; Savira, F.; Suharsono, Y. *Optimization of a Fixed Spraying System for Commercial High-Density Apple Plantings Final Report 2007*; Department of Entomology, Cornell University: Geneva, NY, USA, 2007; Volume 1.
21. Agnello, A.; Landers, A. In-Canopy Fixed Spraying System for High-Density Apple Orchards. In Proceedings of the Communication at Interpoma 2014 Congress, Bolzano, Italy, 20–22 November 2014.
22. Commission of the European Communities Commission regulation (EC) no 889/2008 of 5 September 2008 laying down detailed rules for the implementation of council regulation (EC) no 834/2007 on organic production and labelling of organic products with regard to organic production, labelling and co. *Off. J. Eur. Union* **2008**, *L 250*, 1–84.
23. Townsend, G.K.; Heuberger, J.W. Methods for Estimating Losses Caused by Diseases in Fungicide Experiments. *Plant Dis. Report.* **1943**, *27*, 340–343.
24. Regione del Veneto Decreto 34 del 4 Giugno 2020. Available online: https://veneto.coldiretti.it/wp-content/uploads/2020/06/FD-Decreto-34-del-4_6_2020.pdf (accessed on 28 August 2021).
25. Niemann, S.M.; Whiting, M.D.; Hanrahan, I. Efficacy of a solid set canopy delivery system (SSCDS) in sweet cherry and apple. *Acta Hort.* **2016**, *1130*, 145–150. [[CrossRef](#)]
26. Owen-Smith, P.; Perry, R.; Wise, J.; Jamil, R.Z.R.; Gut, L.; Sundin, G.; Grieshop, M. Spray coverage and pest management efficacy of a solid set canopy delivery system in high density apples. *Pest Manag. Sci.* **2019**, *75*, 3050–3059. [[CrossRef](#)] [[PubMed](#)]
27. Chucho, J.; Thiéry, D. Biology and ecology of the Flavescence dorée vector Scaphoideus titanus: A review. *Agron. Sustain. Dev.* **2014**, *34*, 381–403. [[CrossRef](#)]
28. Tirello, P.; Marchesini, E.; Gherardo, P.; Raniero, D.; Rossetto, F.; Pozzebon, A.; Duso, C. The control of the american leafhopper *Erasmoneura vulnerata* (Fitch) in European vineyards: Impact of synthetic and natural insecticides. *Insects* **2021**, *12*, 85. [[CrossRef](#)]

-
29. Fornasiero, D.; Pavan, F.; Pozzebon, A.; Picotti, P.; Duso, C. Relative infestation level and sensitivity of grapevine cultivars to the leafhopper *Empoasca vitis* (Hemiptera: Cicadellidae). *J. Econ. Entomol.* **2016**, *109*, 416–425. [[CrossRef](#)] [[PubMed](#)]
 30. Duso, C.; Van Leeuwen, T.; Pozzebon, A. Improving the compatibility of pesticides and predatory mites: Recent findings on physiological and ecological selectivity. *Curr. Opin. Insect Sci.* **2020**, *39*, 63–68. [[CrossRef](#)] [[PubMed](#)]